

Comment on <<White-Noise-Induced Transport in Periodic Structures>> by J.Łuczka *et al.*

T. Hondou(*) and Y. Sawada(**)

(*) Graduate School of Information Sciences, Tohoku University,
Sendai, 980-77 Japan

(**) Research Institute of Electrical Communication, Tohoku University
Sendai, 980-77 Japan

PACS. 05.40+j - Fluctuation phenomena, random processes, and Brownian motion.

PACS. 82.20Mj - Nonequilibrium kinetics.

Abstract

In the paper by J.Łuczka *et al.* (*Europhys. Lett.*, **31** (1995) 431), the authors reported by rigorous calculation that an additive Poissonian white shot noise can induce a macroscopic current of a dissipative particle in a periodic potential – even *in the absence* of spatial asymmetry of the potential. We argue that their main result is an obvious one caused by the spatially broken symmetry of a probability distribution of the additive noise, unlike the similar result caused by chaotic noise which has a symmetric probability distribution (*J.Phys.Soc.Jpn.*, **63** (1994) 2014).

In a recent letter [1] in this journal, it was reported that a dissipative particle in a periodic potential can be transported macroscopically by a white shot noise of zero average, even *in the absence* of spatial asymmetry of the potential. Namely, the following dynamics(eq.(1) and eq.(2) of [1] respectively) was considered

$$\dot{x} = -\frac{\partial V}{\partial x} + \xi(t), \quad (1)$$

with

$$\xi(t) = \sum_{i=1}^{n(t)} y_i \delta(t - t_i) - \lambda \langle y_i \rangle, \quad (2)$$

where $V(x)$ is a periodic potential, $\xi(t)$ is a white shot noise, and $n(t)$ is a Poisson counting process with a parameter λ which determines the average sojourn time between two δ -kicks. The positive-valued weights of the δ -pulses $\{y_i\}$, being independent of $n(t)$, are exponentially distributed (eq.(3) of [1]),

$$\rho(y) = A^{-1} \exp[-y/A], \quad y \geq 0, \quad (3)$$

with $\langle y_i \rangle \equiv A$. In this dynamics, the authors of ref.[1] claim that they obtained and analyzed a unidirectional motion of the particle for finite values of the parameters, λ and A , and found an analytical expression for a stationary current of a particle flow.

This subject, noise induced transport, is interesting, in principle, as the authors discussed; however, their main result is trivial from a physical insight, because an existence of a net current is immediately obtained before one performs rigorous calculation, but only in terms of a spatial asymmetry of the additive noise. Their claim that the first term of the additive noise dominates the finite current (see, *r.h.s.* of eq.(2)) is "not evident *a priori*, because the δ -spikes act over zero duration" is not justified as one sees below.

Let us restrict ourselves, for simplicity, in the case of the piecewise linear potential (eq.(17) of ref.[1]). Then, one can find, at a glance, the condition that a particle cannot move at all in a left direction is

$$|\frac{\partial V}{\partial x}| \geq \lambda \langle y_i \rangle, \quad (4)$$

i.e. $1 \geq \lambda A \equiv a_c$. In this condition, however, there exist a finite probability for a particle to move in a right direction, as found in eq.(3): *i.e.*

$$\int_{L/2}^{\infty} \rho(y) dy = \exp[-L/(2A)] > 0 \quad (5)$$

gives a probability that a particle at one of the minima of the potential is transferred to a next basin by a δ -kick of an additive noise at once (see eq.(17) of ref.[1]), where L is a period of the potential. This directly proves an existence of a unidirectional motion of the particle. Therefore, it is evident *a priori* that the particle can be transported unidirectionally for $\lambda A \leq 1$. For a region $\lambda A > 1$, the way how a particle moves is not evident *a priori*; however, an emergence of a net current is not surprising, because the spatial symmetry of the probability distribution of the additive noise is already broken: it would be rather surprising if the net current disappeared at finite parameters, λ and A .

Finally, we remark that a particle can be transported unidirectionally in a similar potential with spatial symmetry even if a probability density of an added noise is *symmetric* and δ correlated[2]. The unidirectional motion can be attributed to a chaotic coherence[3] of which the effect is caused by a *dynamical* asymmetry.

References

- [1] Łuczka J., Bartussek R. and Hänggi P., *Europhys. Lett.*, **31** (1995) 431.
- [2] a) Hondou T., *J.Phys.Soc.Jpn.*, **63** (1994) 2014; b) Hondou T. and Sawada Y., *Phys.Rev.Lett.* **75** (1995) 3269.
- [3] See, for example, Schuster H.G., *Deterministic Chaos* (Physik-Verlag, Weinheim, 1984).